

# Quantitative Analysis of Erythrocytes Containing Fetal Hemoglobin (F Cells) in Children With Sickle Cell Disease

Stacy J. Marcus, Thomas R. Kinney, William H. Schultz, Erin E. O'Branski, and  
Russell E. Ware\*

Department of Pediatrics, Duke-UNC Comprehensive Sickle Cell Center and the Division of Hematology/Oncology,  
Duke University Medical Center, Durham, North Carolina

Variation in the level of fetal hemoglobin (HbF) accounts for much of the clinical heterogeneity observed in patients with sickle cell disease (SCD). The HbF level has emerged as an important prognostic factor in both sickle cell pain and mortality, and a % HbF of 10–20% has been suggested as a threshold level for diminished clinical severity. The number of erythrocytes that contain HbF (termed F cells) may also be critically important, as F cells resist intravascular sickling and have preferential in vivo survival. Since F cells can be enumerated with high accuracy using flow cytometry methods, we prospectively studied a cohort of 242 children with SCD. Children with HbS and hereditary persistence of fetal hemoglobin (S/HPFH) had essentially 100% F cells. In contrast, children with homozygous sickle cell anemia (HbSS), HbS/ $\beta^0$  thalassemia, or HbS/ $\beta^+$  thalassemia had significantly lower mean % F cell values (55.9, 61.6, and 51.3%, respectively;  $P < 0.001$ ), and children with HbSC had even fewer F cells (27.0%;  $P < 0.001$ ). There was a highly significant correlation between the % F cells and the log (% HbF), which was observed for the total population of children ( $r = 0.95$ ,  $P < 0.001$ ), as well as for each of the individual subgroups of children with HbSS ( $r = 0.94$ ,  $P < 0.001$ ), HbSC ( $r = 0.89$ ,  $P < 0.001$ ), or HbS/ $\beta^0$  thalassemia and HbS/ $\beta^+$  thalassemia ( $r = 0.95$ ,  $P < 0.001$ ). This logarithmic correlation between % F cells and % HbF has not been previously described and has important implications for the pharmacologic manipulation of HbF in patients with SCD. *Am. J. Hematol.* 54:40–46, 1997 © 1997 Wiley-Liss, Inc.

**Key words:** children; erythrocytes; F cells; fetal hemoglobin; sickle cell disease

## INTRODUCTION

The clinical manifestations of sickle cell disease (SCD) result from chronic hemolytic anemia and the effects of intravascular sickling, which include local tissue hypoxia and organ damage [1,2]. Patients with SCD commonly develop painful vaso-occlusive events and have an increased risk of infection secondary to splenic infarction. Other common clinical events include acute chest syndrome, splenic sequestration, avascular necrosis, cholelithiasis, priapism, and cerebral vascular accident. There is a large clinical heterogeneity observed in SCD, even for patients with an identical hemoglobin phenotype [3–5]. This variation in clinical severity can be explained in part by differences in the total hemoglobin concentration, the mean corpuscular hemoglobin concentration, erythrocyte rheology, the percentage of adhesive cells, the proportion of dense cells, the presence or absence of  $\alpha$ -thalassemia, and the  $\beta$ -globin haplotype [3,6–9].

The percentage of fetal hemoglobin (% HbF), however, is perhaps the most important laboratory parameter influencing clinical severity in SCD [10,11]. In normal individuals, HbF comprises only 5% of the total hemoglobin by age 3–6 months and falls to below 1% in adults [12]. In contrast, patients with SCD typically have HbF levels ranging from 1 to 20% [13]; those with genetic mutations leading to hereditary persistence of fetal hemoglobin (HPFH) can have HbF levels that reach 30–40% of the

Contract grant sponsors: National Heart, Lung, and Blood Institute, National Institutes of Health and Children's Miracle Network Telethon; Contract grant number: PO 60-HL-28393.

\*Correspondence to: Dr. Russell E. Ware, P.O. Box 2916, Duke University Medical Center, Durham, NC 27710.

Received 4 April 1996; Accepted 15 August 1996

total hemoglobin [14]. Several studies have demonstrated that an increased % HbF is associated with decreased clinical severity in SCD, using endpoints such as the number of painful events, transfusions, and hospitalizations [3,15,16]. Other studies have failed to show a protective effect of HbF, perhaps because the HbF levels were too low to provide protection [7,17]. A potential threshold of 10–20% HbF has been suggested, above which patients experience fewer clinical events [18]. The % HbF has also emerged as the most important predictor of early mortality in patients with SCD [10,19].

Patients with SCD have elevated HbF levels in part due to their increased rate of erythropoiesis [12], but multivariate statistical analysis has documented that age, gender, the presence of  $\alpha$ -thalassemia, the  $\beta$ -globin haplotype, and an X-linked locus also influence the total % HbF within a given individual [5,20–22]. Except in the case of HPFH, HbF is not found in all erythrocytes, but rather is located in a subset known as HbF-containing cells or “F cells” [23,24]. In normal adults, the percentage of F cells ranges from 0.5 to 7%, while in patients with SCD, the % F cells has a much broader range [14,23]. In a few patients with sickle cell anemia tested serially, the % F cells was stable over time within an individual, although the % F cell value varied greatly among different patients [23]. Because F cells have a decreased tendency toward sickle formation, they tend to survive preferentially in the peripheral blood of patients with SCD [23]. The number of F cells, therefore, may be of equal or even greater importance than the absolute amount of HbF in influencing the clinical severity of an affected individual. To date, no prospective analysis of F cells in patients with sickle cell disease has been reported. We therefore quantitated the % F cells in a large cohort of children with SCD and correlated the results with other laboratory parameters. We identified a significant logarithmic correlation between % F cells and % HbF, which has important implications for the pharmacologic increase of HbF in patients with SCD.

## MATERIALS AND METHODS

### Patient Population

All patients received medical care at the Duke Pediatric Hematology Clinic within the Duke-UNC Comprehensive Sickle Cell Center. The diagnosis of a sickle hemoglobinopathy was established using standard hemoglobin electrophoresis techniques, and the % HbF was determined by alkali denaturation. The diagnosis of HbS/ $\beta^+$  thalassemia was established by the presence of HbA (typically 20–25%) in an untransfused sample. The diagnosis of  $\beta^0$  thalassemia was established by the presence of an elevated amount ( $> 3.5\%$ ) of HbA<sub>2</sub> quantitated by column chromatography [25]. When possible, family studies were also performed. All patients in this study were at least 2

years of age, and no patient had been transfused within 4 months.

### F Cell Enumeration

Quantitation of F cell percentage was performed using a slight modification of a previously described protocol [26]. Briefly, whole blood (0.3 mL collected in EDTA) was washed twice with borate saline. To crosslink and fix the erythrocytes, Dimethyl 3,3'-dithiobispropionimidate (DTBP, Pierce, Rockford IL) was added to 1 mL of washed cells, followed by cold borate saline containing diethanolamine. The fixed erythrocytes were resuspended in PBS/2% (phosphate-buffered saline) albumin and stored in 1 mL aliquots at  $-70^\circ\text{C}$  until further analysis.

Fixed erythrocytes were thawed on ice and permeabilized using Triton X-100 and isopropanol, then aliquoted into two samples for immunophenotype analysis. One sample was incubated with a monoclonal antibody (mAb) specific for the  $\gamma$ -globin chain of fetal hemoglobin and has no reactivity with  $\beta$ -globin chains (Immuno-Rx, Atlanta GA). The other sample was incubated with a negative control mAb. After the primary mAb incubation, cells were washed with 0.8% Triton in PBS and then incubated with a goat anti-mouse F(ab')<sub>2</sub> conjugated to fluorescein (Pierce, Rockford IL). The cells were then washed twice and resuspended in 0.5 mL 0.8% Triton in PBS. Five thousand stained cells in each sample were analyzed using a FACSCAN flow cytometer (Beckton-Dickinson) and Lysis II software. Figure 1 shows six examples of F cell enumeration, on cord blood, normal erythrocytes, and samples from four children with sickle cell disease. Each histogram shows a distinct population of cells that stains positively for  $\gamma$ -globin and therefore represents F cells.

### Statistical Analysis


The software program Primer (McGraw-Hill, New York, NY) was used for statistical analysis. Comparison of laboratory parameters with F cell quantitation was performed using linear regression, least squares analysis, ANOVA, and Student's paired *t* test.

## RESULTS

### Patient Population

A total of 242 pediatric patients were included in this study, including 134 males and 108 females. Table I shows that the majority of patients had homozygous sickle cell anemia (HbSS) ( $n = 130$ ), followed by HbSC ( $n = 74$ ), HbS/ $\beta^+$  thalassemia ( $n = 23$ ), HbS/ $\beta^0$  thalassemia ( $n = 7$ ), and HbS/HPFH ( $n = 8$ ). For the entire group of children, the average age was  $8.6 \pm 4.8$  years (mean  $\pm$  1 standard deviation; range, 2.0–19.5 years). There were no significant differences in the ages of children with each hemoglobin genotype (Table I).

The results of quantitative enumeration of % F cells



**Fig. 1. Enumeration of erythrocytes containing HbF (F cells).** Circulating erythrocytes were stained for intracellular  $\gamma$ -globin chains as described in Materials and Methods. Six examples of flow cytometric histograms are shown. A: F cell enumeration in a sample of cord blood, with 96% of cells staining positively for HbF. The isotype control antibody also is shown. B: F cell enumeration on erythrocytes from a normal

donor, with only a small proportion of positive cells. C: A patient with HPFH has >99% F cells. D: A child with HbSS and 23% F cells. E: A child with 51% F cells. F: A child with 79% F cells. In each patient with sickle cell disease, the quantitation of F cells is straightforward, due to the clear separation between erythrocytes that contain fetal hemoglobin and those that do not.


**TABLE I. Clinical and Laboratory Parameters for 242 Children With Sickle Cell Disease\***

Parameter	Hb S/HPFH	Hb SS	Hb S/ $\beta^0$ thalassemia	Hb S/ $\beta^+$ thalassemia	Hb SC
Total patients	8	130	7	23	74
Males	5	73	3	12	40
Females	3	57	4	11	34
Age (years)	$10.1 \pm 7.8$	$8.7 \pm 4.8$	$6.6 \pm 3.6$	$8.4 \pm 4.7$	$8.4 \pm 4.7$
F cells					
%	$98.9 \pm 0.9$	$55.9 \pm 19.9$	$61.6 \pm 27.3$	$51.3 \pm 14.0$	$27.0 \pm 15.8$
Range	96.9 – 99.8	16.7 – 94.4	23.2 – 91.4	29.5 – 85.2	5.1 – 72.7
Hb F					
%	$31.7 \pm 3.0$	$10.4 \pm 6.3$	$15.4 \pm 10.3$	$9.2 \pm 5.0$	$3.3 \pm 2.9$
Range	26.8–37.4	1.5–28.9	3.4–28.0	3.2–21.8	0.5–14.6
Hgb (g/dL)	$12.8 \pm 1.4$	$8.0 \pm 1.0$	$8.4 \pm 1.7$	$10.7 \pm 1.1$	$10.8 \pm 1.1$
Reticulocytes (%)	$1.8 \pm 0.4$	$14.3 \pm 6.4$	$7.0 \pm 2.6$	$2.2 \pm 1.1$	$3.9 \pm 2.5$
Mean corpuscular volume (fL)	$78 \pm 9$	$85 \pm 8$	$69 \pm 4$	$69 \pm 5$	$73 \pm 6$
Mean corpuscular Hgb (pg)	$26.8 \pm 2.6$	$29.0 \pm 3.2$	$22.6 \pm 1.9$	$23.2 \pm 2.1$	$25.6 \pm 2.1$
Mean corpuscular Hgb concentration (g/dL)	$34.7 \pm 1.1$	$34.0 \pm 1.2$	$32.7 \pm 1.2$	$33.7 \pm 0.8$	$35.1 \pm 1.0$
White blood cells ( $\times 10^9/L$ )	$7.7 \pm 2.5$	$12.9 \pm 4.3$	$10.0 \pm 2.8$	$7.8 \pm 2.7$	$8.1 \pm 2.3$
Platelets ( $\times 10^9/L$ )	$299 \pm 138$	$494 \pm 178$	$388 \pm 234$	$326 \pm 105$	$316 \pm 149$

\*Values represent the mean  $\pm$  1 standard deviation for each group.

are summarized in Table I. Experimental values are from a single measurement, although 49 older children with serial measurements demonstrated a high degree of stability over time ( $0.8 \pm 0.7\%$  variability in % HbF,  $4.9 \pm 3.4\%$  variability in % F cells). Eight patients with HbS/HPFH had virtually 100% F cells, which was significantly higher than the mean values for children with

HbSS, HbS/ $\beta^0$  thalassemia, or HbS/ $\beta^+$  thalassemia (55.9, 61.6, and 51.3%, respectively;  $P < 0.001$  for each comparison). There were no significant differences in the mean F cell enumeration among these latter three groups, and their ranges of % F cell values were similar. Children with HbSC had even fewer F cells than any other group of children (27.0%,  $P < 0.001$  for each comparison).



**Fig. 2.** Comparison of % F cells with % HbF for 242 pediatric patients with sickle cell disease, including 130 with HbSS, 74 with HbSC, 23 with HbS/ $\beta^+$  thalassemia, 8 with HbS/HPFH, and 7 with HbS/ $\beta^0$  thalassemia. The F cell enumeration was performed as described in Materials and Methods, and % HbF was quantitated by alkali denaturation. The % F cells was significantly correlated with log (% HbF) ( $r = 0.95, P < 0.001$ ).

Quantitation of fetal hemoglobin revealed a similar pattern (Table I). The children with HbS/HPFH had a mean fetal hemoglobin concentration of 31.7%, which was significantly higher than the mean % HbF for children with HbSS, HbS/ $\beta^0$  thalassemia, or HbS/ $\beta^+$  thalassemia (10.4, 15.4, and 9.2%, respectively;  $P < 0.001$  for each comparison). Children with HbSC had a significantly lower mean % HbF (3.3%;  $P < 0.001$  for each comparison).

Additional hematologic parameters are also listed for each subset of patients, including the hemoglobin concentration, reticulocyte count as determined by an automated technique, red cell indices, white blood cell count, and platelet count (Table I).

#### Comparison of % F Cells With Other Laboratory Parameters


The percentage of F cells was compared with the other laboratory parameters listed in Table I. A highly significant correlation was found between the % F cells and the % HbF. For the entire patient population, the % HbF increased logarithmically with a linear increase in the % F cells (Fig. 2,  $r = 0.95, P < 0.001$ ). This relationship was next analyzed according to hemoglobin genotype.

Within each group, a similar strong correlation was found between % F cells and log (% HbF) (Fig. 3): for the 130 children with HbSS,  $r = 0.94$  ( $P < 0.001$ ); for the 74 children in the HbSC group,  $r = 0.89$  ( $P < 0.001$ ); and for the combined HbS/ $\beta$  thalassemia group (7 with HbS/ $\beta^0$  thalassemia and 23 with HbS/ $\beta^+$  thalassemia),  $r = 0.95$  ( $P < 0.001$ ).

For the children with HbSS, a significant correlation was also found between the % F cell value and the hemoglobin concentration ( $r = 0.54; P < 0.001$ ). For each of the other subgroups, no statistically significant correlation was identified between these two variables.

#### DISCUSSION

The tendency toward intracellular HbS polymerization is probably the most important determinant of SCD pathophysiology [27,28]. Theoretically, therefore, the presence of intracellular HbF is advantageous for several reasons. First, an increased amount of HbF within the erythrocyte decreases the proportion of HbS within that cell. As a consequence, fewer HbS tetramers form, and less hemoglobin polymerization occurs within the cell [29]. In addition, HbF does not co-polymerize well with HbS, thereby



**Fig. 3.** Comparison of % F cells with % HbF according to hemoglobin genotype. **A:** Data for children with HbSS ( $n = 130$ ,  $r = 0.94$ ,  $P < 0.001$ ). **B:** Patients with HbSC ( $n = 74$ ,  $r = 0.89$ ,  $P < 0.001$ ). **C:** Correlation for children with HbS/ $\beta$  thalassemia ( $n = 23$  with HbS/ $\beta^+$  thalassemia and  $n = 7$  with HbS/ $\beta^0$  thalassemia;  $r = 0.95$ ,  $P < 0.001$ ).

increasing the solubility of intracellular HbS [30–33]. Finally, a greater amount of HbF within a cell increases the likelihood of the inclusion of  $\gamma$ -globin chains within the hemoglobin tetramers. These hybrid hemoglobin tetramers hinder HbS polymerization and may be completely excluded from HbS polymers [29,32,34].

Over 30 years ago, Shepard and colleagues [35] used a modification of the Kleihauer-Betke stain to determine that patients with SCD had a variable number of F cells. Analyzing the peripheral blood of eight patients, they noted an unequal distribution of fetal hemoglobin within

the circulating erythrocytes. To date, measurement of the % F cells in large numbers of patients with SCD has not been reported. We used a quantitative assay for F cell enumeration in our study and found that children with HbSS, HbS/ $\beta^0$  thalassemia, or HbS/ $\beta^+$  thalassemia had very similar % F cell values, with a mean of approximately 50–60% F cells for each group and a range of F cell values from approximately 20 to 90%. In contrast, children with HbSC had significantly fewer F cells on average (27.0%) and also a lower mean % HbF (3.3%).

The increase in F cells in SCD patients has been attributed to an increased rate of erythropoiesis [12,14]; this hypothesis is supported by the reports of increased % F cells following recovery from phlebotomy, transient erythroblastopenia [36], or glucose-6-phosphate dehydrogenase (G6PD)-deficient hemolysis [37]. Perhaps children with HbSS have significantly higher % F cells than children with HbSC due to a higher rate of erythropoiesis, as measured by their baseline hemoglobin concentration and reticulocyte counts. However, children with HbS/ $\beta^+$  thalassemia had average hemoglobin and reticulocyte counts very similar to those of the HbSC patients (Table I) but had significantly higher % F cells (51.3% vs. 27.0%;  $P < 0.001$ ). Therefore, the % F cells cannot be based solely on the rate of erythropoiesis. A variety of factors appear to influence the % F cells, including an incompletely understood X-linked F cell production locus [20,22].

Perhaps the most important observation from our analysis was the highly significant logarithmic correlation between the % F cells and the % HbF, which has not been previously described. In an older study, which used a less accurate method of F cell enumeration, Wood et al. [14] found a linear correlation between these two parameters, but the % HbF ranged from only 0 to 3% in their population. Our patients had a much broader range and higher % HbF values, which allowed us to identify the logarithmic correlation (Figs. 2, 3). An important implication of this observation is that the average amount of HbF per F cell varies among individual patients with SCD. For example, if each F cell had a fixed amount of intracellular HbF, then a patient with 60% F cells should have 3 times the % HbF as a patient with only 20% F cells. This would be the result of a linear correlation between the two parameters. The logarithmic correlation is illustrated by the observation that a threefold difference in % F cells (60% vs. 20%) is associated with a fivefold difference in % HbF (10% vs. 2%). This logarithmic correlation was observed for all subsets of SCD patients, but the mechanisms responsible for this highly significant logarithmic association are unknown. It is possible that the % F cells is genetically predetermined in a given patient and that the amount of HbF per F cell is somehow regulated by the total number of F cells within the erythroid progenitor pool.

The logarithmic association between % F cells and % HbF also has important implications for the pharmacologic modification of HbF in patients with SCD. Hydroxyurea is currently being used to increase HbF production in SCD and increases both the % F cells and the % HbF [38–40]. Charache et al. [39] observed a mean increase in HbF of 11% after treatment with hydroxyurea. For a patient starting with 2% HbF, this increase in HbF to 13% would increase the % F cells from approximately 20 to 70%. In contrast, an identical incremental increase in % HbF from 10 to 21% would correspond to an increase in % F cells from 60 to 80%. In terms of increasing the number of F cells that could resist intracellular HbS polymerization, pharmacologic manipulation might be more beneficial for patients who begin with lower % HbF levels. However, patients who begin with higher % HbF levels would have a greater increase in the amount of HbF per F cell, which would also presumably provide a clinical benefit. Quantitative measurement of both % HbF and % F cells should be performed in SCD patients receiving agents such as hydroxyurea and should be correlated with changes in other laboratory or clinical parameters. Greater experience with these pharmacologic agents will determine which patients benefit most.

## ACKNOWLEDGMENTS

The authors thank Mary Whorton for performing quantitative % HbF analysis on these patients, Bethany Bergamo for technical assistance, and Amy Walker for her dedication to the completion of this study. This work was supported in part by grant PO 60-HL-28393 from the National Heart, Lung, and Blood Institute, National Institutes of Health, and by the Children's Miracle Network Telethon.

## REFERENCES

- Platt OS, Dover GJ: Sickle cell disease. In Nathan DG, Oski FA (eds): "Hematology of Infancy and Childhood." Philadelphia: W.B. Saunders, 1993, pp 732–782.
- Charache S, Lubin B, Reid CD (eds): "Management and Therapy of Sickle Cell Disease." 1991.
- Platt OS, Thorington BD, Brambilla DJ, Milner PF, Rosse WF, Vichinsky E, Kinney TR: Pain in sickle cell disease. *N Engl J Med* 325:11, 1991.
- Powars DR:  $\beta^s$ -gene-cluster haplotypes in sickle cell anemia. *Hematol Oncol Clin North Am* 5:475, 1991.
- Steinberg MH, Hsu H, Nagel RL, Milner PF, Adams JG, Benjamin L, Fryd S, Gillette P, Gilman J, Josifovska O, Hellman-Erlingsson S, Safaya S, Huey L, Rieder RF: Gender and haplotype effects upon hematological manifestations of adult sickle cell anemia. *Am J Hematol* 48:175, 1995.
- Steinberg MH, Rosenstock W, Coleman MB, Adams JG, Platiga O, Cedenp M, Rieder RF, Wilson JT, Milner P, West S: The effects of alpha thalassemia and microcytosis on the hematologic and vaso-occlusive severity of sickle cell anemia. *Blood* 63:1353, 1984.
- Baum KF, Dunn DT, Maude GH, Serjeant G: The painful crisis of homozygous sickle cell disease: A study of risk factors. *Arch Intern Med* 147:1231, 1987.
- Powars DR: Sickle cell anemia:  $\beta^s$ -gene-cluster haplotypes as prognostic indicators of vital organ failure. *Semin Hematol* 28:202, 1991.
- Phillips G, Coffey B, Tran-son-tay R, Kinney TR, Orringer EP, Hochmuth RM: Relationship of clinical severity to packed cell rheology in sickle cell anemia. *Blood* 78:2735, 1991.
- Platt OS, Brambilla DJ, Rosse WF, Milner PF, Castro O, Steinberg MH, Klug PP: Mortality in sickle cell disease. *N Engl J Med* 330:1939, 1994.
- Charache S: Fetal hemoglobin, sickling, and sickle cell disease. *Adv Pediatr* 37:1, 1990.
- Wood WG: Increased HbF in adult life. *Baillieres Clin Haematol* 6:177, 1993.
- Serjeant GR: Foetal haemoglobin in homozygous sickle cell disease. *Clin Haematol* 4:109, 1975.
- Wood WG, Stamatoyannopoulos G, Lim G, Nute PE: F-cells in the adult: Normal values and levels in individuals with hereditary and acquired elevations of HbF. *Blood* 46:671, 1975.
- Rucknagel DL, Sarnaik SA, Whitten CF, Odenheimer DA: Fetal hemoglobin concentration predicts disease severity in children with sickle cell anemia. In Stamatoyannopoulos G, Nienhuis AW (eds): "Progress in Clinical and Biological Research, 251: Developmental Control of Globin Gene Expression." New York: Alan R. Liss, 1987, p 487.
- Odenheimer DJ, Sarnaik SA, Whitten CF, Rucknagel DL, Sing CF: The relationship between fetal hemoglobin and disease severity in children with sickle cell anemia. *Am J Med Genet* 27:525, 1987.
- Powars DR, Schroeder WA, Weiss JN, Chan LS, Azen SP: Lack of influence of fetal hemoglobin levels or erythrocyte indices on the severity of sickle cell anemia. *J Clin Invest* 65:732, 1980.
- Powars DR, Weiss JN, Chan LS, Schroeder WA: Is there a threshold of fetal hemoglobin that ameliorates morbidity in sickle cell anemia? *Blood* 63:921, 1984.
- Leikin SL, Gallagher D, Kinney TR, Sloane D, Klug P, Rida W: Mortality in children and adolescents with sickle cell disease. *Pediatrics* 84:500, 1989.
- Miyoshi F, Kaneto Y, Kawai H, Ohchi H, Niki S, Hasegawa K, Shirakami A, Yamano T: X-linked dominant control of F-cells in normal adult life: Characterization of the Swiss type as hereditary persistence of fetal hemoglobin regulated dominantly by gene(s) on X chromosome. *Blood* 72:1854, 1988.
- Falusi AG, Kulozik AE: Relationship of foetal haemoglobin levels and  $\beta^s$  haplotypes in homozygous sickle cell disease. *Eur J Haematol* 45:1, 1990.
- Chang YC, Smith KD, Moore RD, Serjeant G, Dover GJ: An analysis of fetal hemoglobin variation in sickle cell disease: The relative contributions of the X-linked factor,  $\beta$ -globin haplotypes,  $\alpha$ -globin gene number, gender, and age. *Blood* 85:1111, 1995.
- Dover GJ, Boyer SH, Charache S, Heintzelman K: Individual variation in the production and survival of F cells in sickle-cell disease. *N Engl J Med* 299:1428, 1978.
- Boyer SH, Belding TK, Margolet L, Noyes AN: Fetal hemoglobin restriction to a few erythrocytes (F cells) in normal human adults. *Science* 188:361, 1975.
- Kinney TR, Sawtschenko M, Whorton M, Shearin J, Stine C, Hofman L, Safko R, Vitaglione T, Kaufman RE: Techniques' comparison and report of the North Carolina experience. *Pediatrics* 83(suppl):843, 1989.
- Dover GJ, Boyer SH: Fetal hemoglobin-containing cells have the same mean corpuscular hemoglobin as cells without fetal hemoglobin: A reciprocal relationship between gamma- and beta-globin gene expression in normal subjects and in those with high fetal hemoglobin production. *Blood* 69:1109, 1987.
- Brittenham GM, Schechter AN, Noguichi CT: Hemoglobin S polymerization: Primary determinant of the hemolytic and clinical severity of the sickling syndromes. *Blood* 65:183, 1985.
- Noguichi CT, Rodgers GP, Serjeant G, Schechter AN: Levels of fetal

- hemoglobin necessary for treatment of sickle cell disease. *N Engl J Med* 318:96, 1988.
29. Bookchin RM, Nagel RL, Balazs T: Role of hybrid tetramer formation in gelation of haemoglobin S. *Nature* 256:667, 1975.
  30. Singer K, Singer L: The gelling phenomenon of sickle cell hemoglobin: Its biologic and diagnostic significance. *Blood* 8:1008, 1953.
  31. Bertles JF, Rabinowitz R, Dobler J: Hemoglobin interaction: Modification of solid phase composition in the sickling phenomenon. *Science* 169:375, 1970.
  32. Sunshine HR, Hofrichter J, Eaton WA: Gelation of sickle cell hemoglobin in mixtures with normal adult and fetal hemoglobins. *J Mol Biol* 133:435, 1979.
  33. Poillon WN, Kim BC, Rodgers GP, Noguchi CT, Schechter AN: Sparing effect of hemoglobin F and hemoglobin A<sub>2</sub> on the polymerization of hemoglobin S at physiologic ligand saturations. *Proc Natl Acad Sci USA* 90:5039, 1993.
  34. Benesch RE, Edalji R, Benesch R, Kwong S: Solubilization of hemoglobin S by other hemoglobins. *Proc Natl Acad Sci USA* 77:5130, 1988.
  35. Shepard MK, Weatherall DJ, Conley CL: Semi-quantitative estimation of the distribution of fetal hemoglobin in red cell populations. *Bull J Hopkins Hosp* 110:293, 1962.
  36. Papayannopoulou T, Vichinsky E, Stamatoyannopoulos G: Fetal Hb production during acute erythroid expansion. Observations in patients with transient erythroblastopenia and post-phlebotomy. *Br J Haematol* 44:535, 1980.
  37. Schiliro G, Russo A, Romeo MA, Testa R, Musumeci S, Mancuso GR, Russo G: Increase of F cells during acute hemolysis in glucose-6-phosphate dehydrogenase-deficient males. *Pediatr Res* 19:397, 1985.
  38. Goldberg MA, Brugnara C, Dover GJ, Schapira L, Charache S, Bunn HF: Treatment of sickle cell anemia with hydroxyurea and erythropoietin. *N Engl J Med* 323:366, 1990.
  39. Charache S, Dover GJ, Moore RD, Eckert S, Ballas SK, Koshy M, Milner PFA, Orringer EP, Phillips G, Platt OS, Thomas GH: Hydroxyurea: Effects on hemoglobin F production in patients with sickle cell anemia. *Blood* 79:2555, 1992.
  40. Charache S, Terrin ML, Moore RD, Dover GJ, Barton RB, Eckert SV, McMahon RP, Bonds DR: Effect of hydroxyurea on the frequency of painful crises in sickle cell anemia. Investigators of the Multicenter Study of Hydroxyurea in Sickle Cell Anemia. *N Engl J Med* 332:1317, 1995.